


Amathole District Municipality climate change fact sheet

Eastern Cape, South Africa

MUNICIPAL

Introduction


- This fact sheet is part of a series of district municipality fact sheets developed by the Wits GCI and SANBI. The fact sheets present a summary of observed and projected changes in climate over district municipalities in South Africa. They should be used together with the guidelines presented in the cover page.
- Amathole District Municipality covers an area of approximately 23 577 km², with elevation ranging from sea level along the Indian Ocean coastline to 2 332 m above sea level in the Amathole Mountains in the north.
- Climate varies greatly across Amathole, from semi-arid in the west, to wet and subtropical along the Amathole Mountains and northern coast. The district falls in the summer-rainfall region of South Africa.



Observed climate: rainfall (1981–2000)


Mean annual rainfall

Mean annual rainfall ranges from 430 mm over the semi-arid western interior to more than 1 100 mm over parts of the Amathole Mountains and northern coast.



Extreme rainfall days


Mean annual number of extreme rainfall days range from 2 days in the semi-arid western parts to 12 days over the northern coast.



Observed climate: temperature (1981–2000)


Mean annual temperature

Mean annual temperature ranges from 12 °C in the Amathole Mountains to 21 °C along the coastal areas.



Very hot days

Mean annual number of very hot days range from 2 days over the Amathole Mountains to 14 days over the semi-arid western interior.





Observed climate trends (overview)

- Observed decrease in mean annual rainfall (*low confidence*).
- Observed increase in the frequency of extreme rainfall events (*medium confidence*).
- Observed increase in mean annual temperature and warm extremes (*virtually certain*).
- Observed increases in meteorological and agricultural drought (*low confidence*).

Projected future climate change (overview)

- Projected changes in mean annual rainfall are *uncertain* in the near- and mid-future, but there is *high confidence* in decreases in the far-future.
- Projected increase in the frequency of extreme rainfall events (*medium confidence*).
- Projected increase in mean annual temperature and warm extremes (*virtually certain*); decrease in cold extremes (*high confidence*).
- Projected increase in agricultural and meteorological drought in the far-future (*high confidence*).



Projected future climate change (*detailed*)



Near- and mid-future

- Projected decrease in rainfall over the western parts, but increases in the east (*low confidence*).
- Projected increase in extreme rainfall events (*medium confidence*), particularly over the northern mountainous regions.
- Projected increase in temperature and warm extremes (*virtually certain*); decrease in cold extremes (*very likely*).
- Projected increase in agricultural and meteorological drought (*low confidence*).

Far-future

- Projected decrease in rainfall (*likely*) and corresponding increase in agricultural and meteorological drought (*likely*).
- Projected increase in extreme rainfall events (*likely*), particularly over the eastern parts and northern interior regions.
- Projected increase in temperature and warm extremes (*virtually certain*); decrease in cold extremes (*very likely*).

Climate model projections: model agreement and uncertainties



Mean annual rainfall

- Averaged across the district, projected changes in rainfall for the near- and mid-future are *uncertain*.
- Rainfall decreases are projected in the district in the far-future under low mitigation scenarios (*likely*).
- Partially in response to *virtually certain* temperature increases, agricultural drought is to occur more frequently in the near- and mid-future (*low confidence*) and far-future (*likely*).

Mean annual temperature

- Temperature increases averaged across the district in the near-future are *virtually certain* and may be as high as 1.5 °C.
- Under low mitigation, further temperature increases are *virtually certain* and may approach 2.0 °C in the mid-future and 4.0 °C in the far-future.
- Increases in average temperature will be accompanied by increases in warm temperature extremes such as heatwaves and high fire danger days (*virtually certain*).

Citation:

Engelbrecht, F.A., Maviza, A., Steinkopf, J., Vogel, C., Von Maltitz, G., Yose, P. & Barnett, M. 2025. *Sub-national climate change fact sheets for South Africa*. © South African National Biodiversity Institute (SANBI) and University of the Witwatersrand – Global Change Institute (WITS-GCI). DOI: <https://doi.org/10.5281/zenodo.16962181>.

This work is licensed under CC BY-NC-ND 4.0 (Attribution-NonCommercial-NoDerivatives 4.0 International).

<https://creativecommons.org/licenses/by-nc-nd/4.0/deed.en>.

Contact

- Global Change Institute (GCI), University of the Witwatersrand, Johannesburg, South Africa. Website: www.wits.ac.za/gci
- South African National Biodiversity Institute (SANBI). Website: www.sanbi.org